

# *An operational definition of essential biodiversity variables*

**Dirk S. Schmeller, Jean-Baptiste Mihoub, Anne Bowser, Christos Arvanitidis, Mark J. Costello, Miguel Fernandez, Gary N. Geller, et al.**

**Biodiversity and Conservation**


ISSN 0960-3115

Volume 26

Number 12


Biodivers Conserv (2017) 26:2967-2972

DOI 10.1007/s10531-017-1386-9



**Your article is protected by copyright and all rights are held exclusively by Springer Science+Business Media B.V.. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at [link.springer.com](http://link.springer.com)".**

## An operational definition of essential biodiversity variables

Dirk S. Schmeller<sup>1,2</sup>  · Jean-Baptiste Mihoub<sup>1,3</sup> ·  
Anne Bowser<sup>4</sup> · Christos Arvanitidis<sup>5</sup> · Mark J. Costello<sup>6</sup> ·  
Miguel Fernandez<sup>7,8</sup> · Gary N. Geller<sup>9</sup> · Donald Hobern<sup>10</sup> ·  
W. Daniel Kissling<sup>11</sup> · Eugenie Regan<sup>12</sup> · Hannu Saarenmaa<sup>13</sup> ·  
Eren Turak<sup>14,15</sup> · Nick J. B. Isaac<sup>16</sup>

Received: 13 May 2017 / Accepted: 6 June 2017 / Published online: 12 June 2017  
© Springer Science+Business Media B.V. 2017

**Abstract** The concept of essential biodiversity variables (EBVs) was proposed in 2013 to improve harmonization of biodiversity data into meaningful metrics. EBVs were conceived as a small set of variables which collectively capture biodiversity change at multiple spatial scales and within time intervals that are of scientific and management interest. Despite the apparent simplicity of the concept, a plethora of variables that describes not only biodiversity but also any environmental features have been proposed as potential EBV (i.e.

---

Communicated by David Hawksworth.

---

✉ Dirk S. Schmeller

ds@die-schmellers.de

Jean-Baptiste Mihoub

mihoub@mnhn.fr

Anne Bowser

Anne.Bowser@wilsoncenter.org

Christos Arvanitidis

arvanitidis@her.hcmr.gr

Mark J. Costello

m.costello@auckland.ac.nz

Miguel Fernandez

miguel.fernandez@idiv.de

Gary N. Geller

ggeller@geosec.org

Donald Hobern

dhobern@gbif.org

W. Daniel Kissling

wdkissling@gmail.com

Eugenie Regan

eugenie.regan@thebiodiversityconsultancy.com

Hannu Saarenmaa

hannu.saarenmaa@uef.fi

candidate EBV). The proliferation of candidates reflects a lack of clarity on what may constitute a variable that is essential to track biodiversity change, which hampers the operationalization of EBVs and therefore needs to be urgently addressed. Here, we propose that an EBV should be defined as a biological state variable in three key dimensions (time, space, and biological organization) that is critical to accurately document biodiversity change.

**Keywords** Biodiversity monitoring · Conservation policy · Biodiversity change · Essential biodiversity variables · Biological state variables

In 2013, a framework of six classes (Genetic Composition, Species Populations, Species Traits, Community Composition, Ecosystem Structure, Ecosystem Function) of 22 potential essential biodiversity variables (EBVs), inspired by the Essential Climate Variables (ECVs) (Doherty et al. 2009; Bojinski et al. 2014), was proposed (Pereira et al. 2013). The EBV concept aimed to provide an internationally recognized way to monitor essential aspects of biodiversity, such that data from many kinds of sampling programs can be integrated. Thus it would allow comparison of trends in biodiversity from local to national and global scales over time. Operationalizing the EBV concept requires a hierarchy of importance (essentialness) of candidate variables that incorporates their capacity to detect change reliably (Proença et al. 2016). Progress in EBV development has been

---

Eren Turak  
eren.turak@environment.nsw.gov.au

Nick J. B. Isaac  
njbi@ceh.ac.uk

- <sup>1</sup> Department of Conservation Biology, Helmholtz Center for Environmental Research – UFZ, Permoserstrasse 15, 04318 Leipzig, Germany
- <sup>2</sup> Université de Toulouse; UPS, INPT; EcoLab (Laboratoire Ecologie Fonctionnelle et Environnement), Toulouse, France
- <sup>3</sup> Université Pierre et Marie Curie, CESCO, UMR 7204 MNHN-CNRS-UPMC, Paris, France
- <sup>4</sup> Woodrow Wilson International Center for Scholars, 1300 Pennsylvania Ave, NW, Washington, DC, USA
- <sup>5</sup> Institute of Marine Biology, Biotechnology and Aquaculture, Hellenic Centre for Marine Research, Thalassokosmos, Former US Base at Gournes, 71003 Herakleion, Crete, Greece
- <sup>6</sup> Institute of Marine Science, University of Auckland, Auckland 1142, New Zealand
- <sup>7</sup> German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany
- <sup>8</sup> Instituto de Ecología, Universidad Mayor de San Andrés Cota-Cota, Calle 27 Campus Universitario, La Paz, Bolivia
- <sup>9</sup> Group on Earth Observations (GEO), 1211 Geneva, Switzerland
- <sup>10</sup> Global Biodiversity Information Facility Secretariat, Universitetsparken 15, 2100 Copenhagen, Denmark
- <sup>11</sup> Institute for Biodiversity and Ecosystem Dynamics (IBED), University of Amsterdam, P.O. Box 94248, 1090 GE Amsterdam, The Netherlands

hampered by a lack of clarity on what constitutes a biodiversity variable (i.e. candidate EBV). It is urgent to clarify this issue and we argue here that an EBV is a biological state variable that is measurable at particular points in time and space to document biodiversity change.

The EBV concept proposes an integrative view on biodiversity and was conceived as an intermediate layer between primary observational data and derived indicators (Pereira et al. 2013; Brummitt et al. 2016). As a refinement, an EBV can be defined as the harmonized and standardized totality of all biological data across time and space and along a third axis representing the level of biological organization (gene, individual, species, community, ecosystem, aka biological component), altogether framing an EBV cube (Fig. 1). An EBV cube encapsulates a multidimensional view of a specific biodiversity variable and consists of measurements or estimates of essential aspects of biodiversity that support comparison of the state of biodiversity across space and through time. For advancing EBV development, EBVs need to be clearly distinguished from variables describing pressures and responses of biodiversity, such as ecosystem services or disturbance regimes. For instance, the impacts of pressures such as habitat loss or exploitation can be linked to changes in biological variables, such as population abundance, species distribution or habitat structure but the pressures themselves are not biological and thus outside of our definition of an EBV. For the EBV class Ecosystem Function, disturbance regime would not fall into our definition of an EBV as it is not a biological variable, in contrast to the original paper (Pereira et al. 2013). Disturbance regime is rather a natural or anthropogenic driver of change in ecological processes such as succession and regeneration and ultimately of biodiversity states. For the EBV class Ecosystem Structure, EBVs should only cover the biological components, but should not include abiotic variables (chemical composition, slopes, climatic conditions), despite their importance to aid understanding of why an EBV may change in space and over time. This clarification is important and useful in focusing effort directed towards delivering EBVs and in delimiting the data that should be considered suitable for calculating an EBV. It would also allow partitioning the amount of biodiversity data into sub-components that can be realistically and practically addressed in the real world within the frame of existing constraints defined by, e.g., administrative borders, legislative periods, international assessment reporting periods, etc. These real-world constraints may delimit the extent and grain sizes to which EBVs should be matched. This would facilitate the measurement of EBVs anywhere, at any time, in a way that is consistent and generalizable for corresponding components of biodiversity in any biological system. To summarize:

1. EBVs *are* biological state variables: they *are not* pressure (e.g. exploitation), benefit (e.g. ecosystem services), or response (e.g. proportion of habitat in protected areas) variables. EBVs may be combined with pressure, benefit or response variables in subsequent analytical steps (Fig. 1).

---


<sup>12</sup> The Biodiversity Consultancy, 3E King's Parade, Cambridge CB2 1SJ, UK

<sup>13</sup> University of Eastern Finland, Digitalium, Länsikatu 15, 80110 Joensuu, Finland

<sup>14</sup> Australian Museum, 6 College St, Sydney, NSW 2000, Australia

<sup>15</sup> NSW Office on Environment and Heritage, 10 Valentine Avenue, Parramatta 2150, NSW, Australia

<sup>16</sup> Centre for Ecology & Hydrology, Maclean Building, Benson Lane, Crowmarsh Gifford, Wallingford OX10 8BB, UK




**Fig. 1** The process towards operationalizing essential biodiversity variables (EBV). The filling of the EBV cube requires data collection, data mobilization and data processing as a service from and for research (*upper box*). Once the EBV cube has been populated with data, it can facilitate the measurement and modelling of biodiversity change and thereby provides services for science and especially policy (*lower box*). The whole process from raw data to an EBV cube also represents an important service to the scientific community for conducting more resource-efficient analyses of biodiversity change across large spatial extents. This includes quantifying the underlying drivers and pressures of biodiversity change (Mace and Baillie 2007) as well as anticipating and modelling the effect of biodiversity changes on ecosystem services (Oliver et al. 2015). These analyses, with periodic validation, would then feed into global and regional policy processes to explain observed biodiversity change, to improve the forecast of biodiversity change and to produce global assessment reports

2. EBVs are essential because they are selected to capture biodiversity change within a specified range of spatial and temporal grain and extent. They *are not* agnostic to spatial and temporal scale outside of this range.

In general terms, the process to generate an EBV starts with standardizing, harmonizing, and integrating raw biological observations (i.e., primary data) from different sources over space and time and ends when its spatiotemporal changes are fully documented. The development of indicators and the understanding of the causes of the documented change do not fall within the EBV framework, but are a logical next step in using the EBV data (Figs. 1, 2). Taken in combination, EBVs would be a representative set of critical components of biodiversity (Schmeller et al. in press) and thus would help to prioritize data





**Fig. 2** Examples of potential EBVs, their representation in space for a given time step  $t_x$ , and how they might be used to document temporal changes in biodiversity when comparing different spatial representation along the temporal dimension. For example, counts of individuals at time  $t_x$  constitute one time slice of the EBV species abundance. Once several of these time slices are available, temporal change in species abundance across space can be documented. Analyses combining available data on drivers and pressures and the observed change can then inform the policy sphere on the underlying causes of that change (Fig. 1)

collection methods, management and publication. The final EBV suite would need to be evaluated based on the added benefit (essentialness) of each EBV to all others using a set of objective criteria such as representativity, complementarity, policy relevance, predictive ability, significance of change, sensitivity to change, and applicability across biological realms. The essentialness of an EBV then could be used to prioritise on which EBV to focus efforts on for the measurement and operationalization across sampling programs. It is also important to recognize that different stakeholders and groups of scientists are likely to hold differing views on the suite of candidate EBVs that should be operationalized: policy-makers might favour a narrow set of EBVs that produces unambiguous signals of change based on streamlined infrastructure (data flow pipelines etc.). Scientists have an important role in this debate, to ensure that the final set captures the critical variables to understand the state of and change in biodiversity (Schmeller et al. in press).

In conclusion, our refinement of the EBV concept is an important step towards the operationalization of EBVs. Providing a clear frame what an EBV is (or should be) by its *intrinsic nature* is at least as important for operationalizing it than developing technical, analytical, legal ways to produce an EBV along a workflow. Otherwise, any environmental variable would be included in the EBV concept, weakening the potential impact of the concept considerably. A weakened concept would threaten the potential of EBVs to empower and coordinate biodiversity monitoring globally (Schmeller et al. 2015), to

provide an integrative view of biodiversity and delivering stakeholders with clear and simple information about biodiversity change (Brummitt et al. 2016). Along with our suggestion to frame EBVs as strict biological state variables, we urge biodiversity scientists to embrace a more transparent definition of biodiversity variables, be it similar or different to our proposition. We also encourage scientists to develop complementary essential environmental variables that would capture other critical aspects of our changing planet that are not directly related to biodiversity, such as the valuation of ecosystem services or the chemical composition of soil and water. Once implemented, the EBV framework would allow quantification of past trends and providing key data for forecasting future changes in biodiversity so as to inform meaningful, policy relevant and comprehensive indicators and projections.

**Acknowledgements** The authors would like to thank A. Loyau for an insightful outsider's view.

**Funding** DSS, ER, CA, and JBM were financed by the EU BON project that is a 7th Framework Programme funded by the European Union under Contract No. 308454. CA additionally received funding from the LifeWatch Greece infrastructure (MIS 384676), funded by the Greek Government under the General Secretariat of Research and Technology (GSRT), ESFRI Projects, and National Strategic Reference Framework (NSRF). MF was funded by the GEO-BON Secretariat. AB's participation is supported by the Alfred P. Sloan Foundation. WDK acknowledges support from the European Union's Horizon 2020 research and innovation programme (No. 654003) for the GLOBIS-B project ([www.globis-b.eu](http://www.globis-b.eu)).

**Author contributions** All authors have intensively discussed the refinement of the EBV concept and have contributed to writing the manuscript under the lead of the first and senior author.

## References

- Bojinski S, Verstraete M, Peterson TC et al (2014) The concept of essential climate variables in support of climate research, applications, and policy. *Bull Am Meteorol Soc* 95:1431–1443
- Brummitt N, Regan E, Weatherdon LV et al (2016) Taking stock of nature: essential biodiversity variables explained. *Biol Conserv*. doi:[10.1016/j.biocon.2016.09.006](https://doi.org/10.1016/j.biocon.2016.09.006)
- Doherty RM, Heal MR, Wilkinson P et al (2009) Current and future climate-and air pollution-mediated impacts on human health. *Environ Health* 8:S8
- Mace GM, Baillie JE (2007) The 2010 biodiversity indicators: challenges for science and policy. *Conserv Biol* 21:1406–1413
- Oliver TH, Heard MS, Isaac NJ et al (2015) Biodiversity and resilience of ecosystem functions. *Trends Ecol Evol* 30:673–684
- Pereira HM, Ferrier S, Walters M et al (2013) Essential biodiversity variables. *Science* 339:277–278
- Proença V, Martin LJ, Pereira HM et al (2016) Global biodiversity monitoring: from data sources to essential biodiversity variables. *Biol Conserv*. doi:[10.1016/j.biocon.2016.07.014](https://doi.org/10.1016/j.biocon.2016.07.014)
- Schmeller DS, Juillard R, Bellingham PJ et al (2015) Towards a global terrestrial species monitoring program. *J Nat Conserv* 25:51–57
- Schmeller DS, Weatherdon LV, Loyau A et al (2017) A suite of essential biodiversity variables for detecting critical biodiversity change. *Biol Rev*. doi:[10.1111/brv.12332](https://doi.org/10.1111/brv.12332)